

From Möbius To Gyrogroups

Abraham A. Ungar

Amer. Math. Monthly, Vol. 115(2), (2008), pp. 138–144.

1. INTRODUCTION

The evolution from Möbius to gyrogroups began in 1988 [12], and is still ongoing in [14, 15]. Gyrogroups, a natural generalization of groups, lay a fruitful bridge between nonassociative algebra and hyperbolic geometry, just as groups lay a fruitful bridge between associative algebra and Euclidean geometry. More than 150 years have passed since the German mathematician August Ferdinand Möbius (1790–1868) first studied the transformations that now bear his name [9, p. 71]. Yet, the rich structure he thereby exposed is still far from being exhausted, as the evolution from Möbius to gyrogroups demonstrates.

2. FROM MÖBIUS

Möbius transformations of the complex open unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ of the complex plane $\mathbb{C}$ are studied in most books on function theory of one complex variable. According to [10, p. 176], these are important for at least two reasons: (i) they play a central role in non-Euclidean geometry [15], and (ii) they are the only automorphisms of the disc. Ahlfors' book [1], *Conformal Invariants: Topics in Geometric Function Theory*, begins with a presentation of the *polar decomposition* of the most general Möbius self-transformation of the complex open unit disc $\mathbb{D}$,

$$(1) \quad z \mapsto e^{i\theta} \frac{a+z}{1+\bar{a}z} = e^{i\theta}(a \oplus z),$$

where $a, z \in \mathbb{D}$ and $\theta \in \mathbb{R}$ [3, p. 211] [6, p. 185] [10, pp. 177–178]. We present the Möbius polar decomposition (1) in a form that suggests the *Möbius addition*, $\oplus$, defined by the equation

$$(2) \quad a \oplus z = \frac{a+z}{1+\bar{a}z}.$$

Naturally, Möbius subtraction, $\ominus$, is given by $a \ominus z = a \oplus (-z)$, so that $z \ominus z = 0$ and $\ominus z = 0 \ominus z = 0 \oplus (-z) = -z$.

Interestingly, Möbius addition possesses the *automorphic inverse property*

$$(3) \quad \ominus(a \oplus b) = \ominus a \ominus b$$

and the *left cancellation law*

$$(4) \quad \ominus a \oplus (a \oplus z) = z$$

for all $a, b, z \in \mathbb{D}$.

Möbius addition is neither commutative nor associative. To make it look like a commutative binary operation in the disc, we define the *gyrations*

$$(5) \quad \text{gyr}[a, b] = \frac{a \oplus b}{b \oplus a} = \frac{1 + a\bar{b}}{1 + \bar{a}b}$$

for $a, b \in \mathbb{D}$. Being unimodular complex numbers, gyrations represent rotations of the disc about its center.

Moreover, gyrations respect Möbius addition:

$$(6) \quad \text{gyr}[a, b](c \oplus d) = \text{gyr}[a, b]c \oplus \text{gyr}[a, b]d$$

for all $a, b, c, d \in \mathbb{D}$. Also, the inverse of a gyration $\text{gyr}[a, b]$ is the gyration $\text{gyr}[b, a]$:

$$(7) \quad \text{gyr}^{-1}[a, b] = \text{gyr}[b, a].$$

Hence, gyrations are special automorphisms of the Möbius groupoid $(\mathbb{D}, \oplus)$. We recall that a groupoid $(G, \oplus)$ is a nonempty set, G , with a binary operation, $\oplus$, and that an automorphism of a groupoid $(G, \oplus)$ is a bijective self map f of G that respects its binary operation $\oplus$, that is, $f(a \oplus b) = f(a) \oplus f(b)$. The set of all automorphisms of a groupoid $(G, \oplus)$ forms a group, denoted $\text{Aut}(G, \oplus)$.

To emphasize that gyrations are automorphisms, they are also called *gyroautomorphisms*. The automorphism $\phi(z) = -z$ of the disc is not a gyroautomorphism of the disc since the equation $(1 + a\bar{b})/(1 + \bar{a}b) = -1$ for $a, b \in \mathbb{D}$ has no solution in the disc. Consequently, the gyroautomorphisms of the disc do not form a group under gyroautomorphism composition.

It follows from the gyration definition in (5) that Möbius addition obeys the *gyrocommutative law*:

$$(8) \quad a \oplus b = \text{gyr}[a, b](b \oplus a)$$

for all $a, b \in \mathbb{D}$. The Möbius gyrocommutative law (8) is not terribly surprising since it is generated by the gyration definition. But, we are not finished!

“Coincidentally,” the gyration $\text{gyr}[a, b]$ that “repairs” in (8) the breakdown of commutativity in Möbius addition “repairs” the breakdown of associativity as well, giving rise to the following gyroassociative laws (left and right):

$$(9) \quad \begin{aligned} a \oplus (b \oplus z) &= (a \oplus b) \oplus \text{gyr}[a, b]z, \\ (a \oplus b) \oplus z &= a \oplus (b \oplus \text{gyr}[b, a]z) \end{aligned}$$

for all $a, b, z \in \mathbb{D}$. Hence, gyrations measure simultaneously the extent to which Möbius addition deviates from both associativity and commutativity.

Möbius gyrations are expressible in terms of Möbius addition. Indeed, solving the first identity in (9) for $\text{gyr}[a, b]z$ by means of the Möbius left cancellation law (4) we have

$$(10) \quad \text{gyr}[a, b]z = \ominus(a \oplus b) \oplus \{a \oplus (b \oplus z)\}.$$

Möbius gyrations $\text{gyr}[a, b]$, $a, b \in \mathbb{D}$, thus endow the Möbius disc groupoid $(\mathbb{D}, \oplus)$ with a group-like structure, naturally called a *gyrogroup*. In fact, Möbius gyrations do more than that. They contribute their own rich structure to the gyrogroup. Thus, for instance, they possess the *loop properties* (left and right)

$$(11) \quad \begin{aligned} \text{gyr}[a \oplus b, b] &= \text{gyr}[a, b], \\ \text{gyr}[a, b \oplus a] &= \text{gyr}[a, b], \end{aligned}$$

and the nested gyration identity

$$(12) \quad \text{gyr}[b, \ominus \text{gyr}[b, a]a] = \text{gyr}[a, b].$$

More gyration identities, along with their applications in hyperbolic geometry and Einstein's special relativity, are presented in [14, 15]. An important hyperbolic geometric interpretation of Möbius gyrations as triangle defects is presented in [20], and an important special relativistic interpretation of gyrations as *Thomas precessions* is presented in [16] and noted in [19].

Guided by the group-like structure of Möbius addition, we define a *gyrogroup* to be a group in which the associative law is replaced by the gyroassociative laws (left and right), and a *gyrocommutative gyrogroup* to be a commutative group in which the commutative and associative laws are replaced by the gyrocommutative and gyroassociative laws. Like Möbius gyrations, the gyrations in the gyrocommutative and gyroassociative laws of an abstract gyrogroup must be automorphisms of the gyrogroup.

Thus, the disc $\mathbb{D}$ with its Möbius addition forms a *gyrocommutative gyrogroup* $(\mathbb{D}, \oplus)$ where, strikingly, the concepts of gyrocommutativity and gyroassociativity of the gyrogroup preserve the flavor of their classical counterparts [14, 15]. Mathematical coincidences are not accidental! Hence, in particular, the gyrations that “coincidentally” repair in (8)–(9) the breakdown of commutativity and associativity need not be peculiar to Möbius addition, as shown in [4, 5].

3. TO GYROGROUPS

Taking the key features of the Möbius groupoid $(\mathbb{D}, \oplus)$ as axioms, we are now in a position to present the formal gyrogroup definition.

Definition 1. (Gyrogroups). *A groupoid $(G, \oplus)$ is a gyrogroup if its binary operation satisfies the following axioms. In G there is at least one element, 0 , called a left identity, satisfying*

$$(G1) \quad 0 \oplus a = a$$

for all $a \in G$. There is an element $0 \in G$ satisfying axiom (G1) such that for each $a \in G$ there is an element $\ominus a \in G$, called a left inverse of a , satisfying

$$(G2) \quad \ominus a \oplus a = 0.$$

Moreover, for any $a, b, c \in G$ there exists a unique element $\text{gyr}[a, b]c \in G$ such that the binary operation obeys the left gyroassociative law

$$(G3) \quad a \oplus (b \oplus c) = (a \oplus b) \oplus \text{gyr}[a, b]c.$$

The map $\text{gyr}[a, b] : G \rightarrow G$ given by $c \mapsto \text{gyr}[a, b]c$ is an automorphism of the groupoid $(G, \oplus)$, that is,

$$(G4) \quad \text{gyr}[a, b] \in \text{Aut}(G, \oplus),$$

and the automorphism $\text{gyr}[a, b]$ of G is called the gyroautomorphism, or the gyration, of G generated by $a, b \in G$. The operator $\text{gyr} : G \times G \rightarrow \text{Aut}(G, \oplus)$ is called the gyrator of G . Finally, the gyroautomorphism $\text{gyr}[a, b]$ generated by any $a, b \in G$ possesses the left loop property

$$(G5) \quad \text{gyr}[a, b] = \text{gyr}[a \oplus b, b].$$

The gyrogroup axioms (G1)–(G5) in Definition 1 are classified into three classes:

- (1) The first pair of axioms, (G1) and (G2), is a reminiscent of the group axioms.
- (2) The last pair of axioms, (G4) and (G5), presents the gyrator axioms.
- (3) The middle axiom, (G3), is a hybrid axiom linking the two pairs of axioms in (1) and (2).

As in group theory, we use the notation $a \ominus b = a \oplus (\ominus b)$ in gyrogroup theory as well.

In full analogy with groups, gyrogroups are classified into gyrocommutative and non-gyrocommutative gyrogroups.

Definition 2. (Gyrocommutative Gyrogroups). *A gyrogroup $(G, \oplus)$ is gyrocommutative if its binary operation obeys the gyrocommutative law*

$$(G6) \quad a \oplus b = \text{gyr}[a, b](b \oplus a)$$

for all $a, b \in G$.

Clearly, a (commutative) group is a degenerate (gyrocommutative) gyrogroup whose gyroautomorphisms are all trivial. The algebraic structure of gyrogroups is, accordingly, richer than that of groups. Thus, without losing the flavor of the group structure we have generalized it into the gyrogroup structure to suit the needs of Möbius addition in the disc. Fortunately, the gyrogroup structure is by no means restricted to Möbius addition in the disc. Rather, it abounds in group theory as demonstrated, for instance, in [4] and [5], where finite and infinite gyrogroups, both gyrocommutative and non-gyrocommutative, are studied. Some first gyrogroup theorems, some of which are analogous to group theorems, are presented in [15, Chap. 2].

4. MÖBIUS GYROGROUPS: FROM THE DISC TO THE BALL

The gyrocommutative gyrogroup structure in Definition 2 is tailor-made for Möbius addition in the disc. However, it suits Möbius addition in the ball of any real inner product space as well.

Let us identify complex numbers of the complex plane $\mathbb{C}$ with vectors of the Euclidean plane $\mathbb{R}^2$ in the usual way:

$$(13) \quad \mathbb{C} \ni u = u_1 + iu_2 = (u_1, u_2) = \mathbf{u} \in \mathbb{R}^2.$$

Then, the equations

$$(14) \quad \begin{aligned} \bar{u}v + u\bar{v} &= 2\mathbf{u} \cdot \mathbf{v}, \\ |u| &= \|\mathbf{u}\| \end{aligned}$$

give the inner product and the norm in $\mathbb{R}^2$, so that Möbius addition in the disc $\mathbb{D}$ of $\mathbb{C}$ becomes Möbius addition in the disc $\mathbb{R}_{s=1}^2 = \{\mathbf{v} \in \mathbb{R}^2 : \|\mathbf{v}\| < s = 1\}$ of $\mathbb{R}^2$.

Indeed, it follows from (14) that

$$\begin{aligned}
 \mathbb{D} \ni u \oplus v &= \frac{u+v}{1+\bar{u}v} \\
 &= \frac{(1+u\bar{v})(u+v)}{(1+\bar{u}v)(1+u\bar{v})} \\
 (15) \quad &= \frac{(1+\bar{u}v+u\bar{v}+|v|^2)u + (1-|u|^2)v}{1+\bar{u}v+u\bar{v}+|u|^2|v|^2} \\
 &= \frac{(1+2\mathbf{u}\cdot\mathbf{v}+\|\mathbf{v}\|^2)\mathbf{u} + (1-\|\mathbf{u}\|^2)\mathbf{v}}{1+2\mathbf{u}\cdot\mathbf{v}+\|\mathbf{u}\|^2\|\mathbf{v}\|^2} \\
 &= \mathbf{u} \oplus \mathbf{v} \in \mathbb{R}_{s=1}^2
 \end{aligned}$$

for all $u, v \in \mathbb{D}$ and all $\mathbf{u}, \mathbf{v} \in \mathbb{R}_{s=1}^2$. The last equation in (15) is a vector equation, so that its restriction to the ball of the Euclidean two-dimensional space is a mere artifact. As such, it survives unimpaired in higher dimensions, suggesting the following definition of Möbius addition in the ball of any real inner product space.

Definition 3. (Möbius Addition in the Ball). Let $\mathbb{V}$ be a real inner product space [8], and let $\mathbb{V}_s$ be the s -ball of $\mathbb{V}$,

$$(16) \quad \mathbb{V}_s = \{\mathbf{v}_s \in \mathbb{V} : \|\mathbf{v}\| < s\},$$

for any fixed $s > 0$. Möbius addition $\oplus$ is a binary operation in $\mathbb{V}_s$ given by the equation

$$(17) \quad \mathbf{u} \oplus \mathbf{v} = \frac{(1 + \frac{2}{s^2} \mathbf{u} \cdot \mathbf{v} + \frac{1}{s^2} \|\mathbf{v}\|^2) \mathbf{u} + (1 - \frac{1}{s^2} \|\mathbf{u}\|^2) \mathbf{v}}{1 + \frac{2}{s^2} \mathbf{u} \cdot \mathbf{v} + \frac{1}{s^4} \|\mathbf{u}\|^2 \|\mathbf{v}\|^2},$$

where $\cdot$ and $\|\cdot\|$ are the inner product and norm that the ball $\mathbb{V}_s$ inherits from its space $\mathbb{V}$ and where, ambiguously, $+$ denotes both addition of real numbers on the real line and addition of vectors in $\mathbb{V}$.

Without loss of generality, one may select $s = 1$ in Definition 3. We, however, prefer to keep s as a free positive parameter in order to exhibit the result that in the limit as $s \rightarrow \infty$, the ball $\mathbb{V}_s$ expands to the whole of its real inner product space $\mathbb{V}$, and Möbius addition $\oplus$ reduces to vector addition in $\mathbb{V}$.

Being special automorphisms, Möbius gyrations form an important ingredient of the Möbius disc gyrogroup $(\mathbb{D}, \oplus)$. Following the extension of Möbius addition from the disc to the ball, it is desirable to extend Möbius gyrations from the disc to the ball as well. On first glance this task seems impossible, since Möbius gyrations in the disc are given by (5) in terms of a division of complex numbers by nonzero complex numbers, an operation that cannot be extended from the disc to the ball. Fortunately, however, identity (10) comes to the rescue. This identity expresses gyrations in terms of Möbius addition in the disc, an operation that we have just extended from the disc to the ball in (15), as formalized in Definition 3.

As suggested by (10), and in agreement with Definition 1 of gyrogroups, the definition of Möbius gyrations in the ball follows.

Definition 4. (Möbius Gyrations in the Ball). Let $(\mathbb{V}_s, \oplus)$ be a Möbius groupoid. The gyrator gyr is the map $\text{gyr} : \mathbb{V}_s \times \mathbb{V}_s \rightarrow \text{Aut}(\mathbb{V}_s, \oplus)$ given by

the equation

$$(18) \quad \text{gyr}[\mathbf{a}, \mathbf{b}]\mathbf{z} = \ominus(\mathbf{a} \oplus \mathbf{b}) \oplus \{\mathbf{a} \oplus (\mathbf{b} \oplus \mathbf{z})\}$$

for all $\mathbf{a}, \mathbf{b}, \mathbf{z} \in \mathbb{V}_s$. The automorphisms $\text{gyr}[\mathbf{a}, \mathbf{b}]$ of $\mathbb{V}_s$ are called gyrations, or gyroautomorphisms, of the ball $\mathbb{V}_s$.

It is anticipated in Definition 4 that gyrations of the ball are automorphisms of the ball groupoid $(\mathbb{V}_s, \oplus)$. To show that this is indeed the case, we note that (18) can be manipulated by means of Definition 3 of Möbius addition, obtaining (with the help of computer algebra)

$$(19) \quad \begin{aligned} \text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{w} &= \ominus(\mathbf{u} \oplus \mathbf{v}) \oplus \{\mathbf{u} \oplus (\mathbf{v} \oplus \mathbf{w})\} \\ &= \mathbf{w} + 2 \frac{A\mathbf{u} + B\mathbf{v}}{D}, \end{aligned}$$

where

$$(20) \quad \begin{aligned} A &= -\frac{1}{s^4} \mathbf{u} \cdot \mathbf{w} \|\mathbf{v}\|^2 + \frac{1}{s^2} \mathbf{v} \cdot \mathbf{w} + \frac{2}{s^4} (\mathbf{u} \cdot \mathbf{v})(\mathbf{v} \cdot \mathbf{w}), \\ B &= -\frac{1}{s^4} \mathbf{v} \cdot \mathbf{w} \|\mathbf{u}\|^2 - \frac{1}{s^2} \mathbf{u} \cdot \mathbf{w}, \\ D &= 1 + \frac{2}{s^2} \mathbf{u} \cdot \mathbf{v} + \frac{1}{s^4} \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 \end{aligned}$$

for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{V}_s$. Owing to the Cauchy-Schwarz inequality [8, p. 20], $D > 0$ for $\mathbf{u}$ and $\mathbf{v}$ in the ball $\mathbb{V}_s$. By expanding the domain of $\mathbf{w}$ from the ball $\mathbb{V}_s$ to the space $\mathbb{V}$ in (19)–(20), we can extend the gyrations $\text{gyr}[\mathbf{u}, \mathbf{v}]$ to linear maps of $\mathbb{V}$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}_s$.

It follows from (19)–(20) straightforwardly (the use of computer algebra is recommended) that

$$(21) \quad \text{gyr}[\mathbf{v}, \mathbf{u}](\text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{w}) = \mathbf{w}$$

for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{V}_s$, so that gyrations of the ball are invertible, the inverse of $\text{gyr}[\mathbf{u}, \mathbf{v}]$ being $\text{gyr}[\mathbf{v}, \mathbf{u}]$.

Furthermore, it follows from (19)–(20) straightforwardly (the use of computer algebra is recommended) that Möbius gyrations of the ball preserve the inner product that the ball $\mathbb{V}_s$ inherits from its real inner product space $\mathbb{V}$, that is,

$$(22) \quad \text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{a} \cdot \text{gyr}[\mathbf{u}, \mathbf{v}]\mathbf{b} = \mathbf{a} \cdot \mathbf{b}$$

for all $\mathbf{a}, \mathbf{b}, \mathbf{u}, \mathbf{v} \in \mathbb{V}_s$. This implies, in turn, that Möbius gyrations keep invariant the norm that the ball $\mathbb{V}_s$ inherits from its real inner product space $\mathbb{V}$. As such, Möbius gyrations of the ball preserve Möbius addition in the ball, so that Möbius gyrations are automorphisms of the Möbius ball groupoid $(\mathbb{V}_s, \oplus)$. Being special automorphisms, Möbius gyrations are also called gyroautomorphisms of the ball. The automorphism $\phi(\mathbf{z}) = -\mathbf{z}$ of the ball is not a gyroautomorphism of the ball, as verified in [15, p. 70]. Hence, the gyroautomorphisms of the ball do not form a group under gyroautomorphism composition.

Not unexpectedly, Möbius addition $\oplus$ in the ball $\mathbb{V}_s$ of any real inner product space $\mathbb{V}$ preserves the structure it has in the disc. It thus gives rise to the Möbius gyrocommutative gyrogroup $(\mathbb{V}_s, \oplus)$ in the ball, studied in [15]. Möbius addition in the ball is known in the literature as a *hyperbolic translation* [2, 11]. However, its gyrogroup structure went unnoticed until it was uncovered in 1988 [12, 13] in the context of Einstein's special theory of relativity. Indeed, like Möbius addition, Einstein's velocity addition law of special relativity gives rise to the Einstein gyrocommutative gyrogroup $(\mathbb{V}_s, \oplus_E)$ in the ball, studied in [12, 14, 16].

The evolution of Möbius addition and gyrations in the disc does not stop at the level of gyrogroups. It continues into the regime of *gyrovectors spaces*. Strikingly, gyrovectors spaces form the setting for hyperbolic geometry just as vector spaces form the setting for Euclidean geometry. In particular, Möbius gyrovectors spaces form the setting for the Poincaré ball model of hyperbolic geometry while, similarly, Einstein gyrovectors spaces form the setting for the Beltrami-Klein ball model of hyperbolic geometry [14, 15, 17, 18, 21]. Accordingly, a gyrovectors space approach to analytic hyperbolic geometry, fully analogous to the common vector space approach to Euclidean geometry [7], is developed in [15].

REFERENCES

1. Lars V. Ahlfors. *Conformal invariants: topics in geometric function theory*. McGraw-Hill Book Co., New York, 1973. McGraw-Hill Series in Higher Mathematics.
2. Lars V. Ahlfors. *Möbius transformations in several dimensions*. University of Minnesota School of Mathematics, Minneapolis, Minn. 1981.
3. Stephen D. Fisher. *Complex variables*. Dover Publications Inc. Mineola, NY, 1999. Corrected reprint of the second (1990) edition.
4. Tuval Foguel and Abraham A. Ungar. Involutory decomposition of groups into twisted subgroups and subgroups. *J. Group Theory*, **3** (2000) 27–46.
5. Tuval Foguel and Abraham A. Ungar. Gyrogroups and the decomposition of groups into twisted subgroups and subgroups. *Pac. J. Math.* **197** (2001) 1–11.
6. Robert E. Greene and Steven G. Krantz. *Function theory of one complex variable*. John Wiley & Sons Inc. New York, 1997.
7. Melvin Hausner. *A vector space approach to geometry*. Dover Publications Inc., Mineola, NY, 1998. Reprint of the 1965 original.
8. Jerrold E. Marsden. *Elementary classical analysis*. W. H. Freeman and Co., San Francisco, 1974. With the assistance of Michael Buchner, Amy Erickson, Adam Hausknecht, Dennis Heifetz, Janet Macrae and William Wilson, and with contributions by Paul Chernoff, István Fáry and Robert Gulliver.
9. David Mumford, Caroline Series, and David Wright. *Indra's pearls: The vision of Felix Klein*. Cambridge University Press, New York, 2002.
10. Tristan Needham. *Visual complex analysis*. The Clarendon Press Oxford University Press, New York, 1997.
11. John G. Ratcliffe. *Foundations of hyperbolic manifolds*, volume 149 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
12. Abraham A. Ungar. Thomas rotation and the parametrization of the Lorentz transformation group. *Found. Phys. Lett.* **1** (1988) 57–89.
13. Abraham A. Ungar. Möbius transformations of the ball, Ahlfors' rotation and gyrovectors spaces. In *Themistocles M. Rassias (ed.): Nonlinear analysis in geometry and topology*, pages 241–287. Hadronic Press, Palm Harbor, FL, 2000.
14. Abraham A. Ungar. *Beyond the Einstein addition law and its gyroscopic Thomas precession: The theory of gyrogroups and gyrovectors spaces*, volume 117 of *Fundamental Theories of Physics*. Kluwer Academic Publishers Group, Dordrecht, 2001.

15. Abraham A. Ungar. *Analytic hyperbolic geometry: Mathematical foundations and applications*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005.
16. Abraham A. Ungar. Einstein's special relativity: Unleashing the power of its hyperbolic geometry. *Comput. Math. Appl.* **49** (2005) 187–221.
17. Abraham A. Ungar. Gyrovector spaces and their differential geometry. *Nonlinear Funct. Anal. Appl.* **10** (2005) 791–834.
18. Abraham A. Ungar. The relativistic hyperbolic parallelogram law. In *Geometry, integrability and quantization*. Softex, Sofia, (2006) 249–264.
19. Abraham A. Ungar. Thomas precession: a kinematic effect of the algebra of Einstein's velocity addition law. Comments on 'deriving relativistic momentum and energy: II. three dimensional case'. *European J. Phys.* **27** (2006) L17–L20.
20. J. Vermeer. A geometric interpretation of Ungar's addition and of gyration in the hyperbolic plane. *Topology Appl.* **152** (2005) 226–242.
21. Scott Walter. Book Review: *Beyond the Einstein Addition Law and its Gyroscopic Thomas Precession: The Theory of Gyrogroups and Gyrovector Spaces*, by Abraham A. Ungar. *Found. Phys.* **32** (2002) 327–330.